	1
- 2	7
	•

PTO/SB/05 (4/98)

Approved for use through 9/30/2000. OMB 0651-0032 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Type a plus sign (+) inside this box +

Attorney Docket No. M 6560 OS/OAPT First Inventor or Application Identifier Anderson et al. Title IMPROVED FERMENTATION PROCESS

Express Mail Label No. EL544993025US

UTILITY PATENT APPLICATION **TRANSMITTAL**

Only for new nonprovisional app	lications under 37 C.F.R. § 1.53(b))							, ,
APPLICAT	ION ELEMENTS	ADI	ORE	SS TO:	Box Patent	ommissione Application	r for Patents	
See MPEP chapter 600 concernir	g utility patent application contents.				Washington	n, DC 20231		
* Fee Transmitta (Submit an original and	Form (e.g., PTO/SB/17) a duplicate for fee processing)	5.		Microfiche	Computer P	rogram (App	endix)	
Specification (preferred arrangement)	[Total Pages] 26	6. N	ıcled	tide and/or	Amino Acid 8	Sequence S	ubmission	- 1
- Descriptive title		a.		bie, all necessa Comput	er Readable	Сору		- 1
- Cross Reference	es to Related Applications	b.	г	Paper C	copy (identica	I to compute	er copy)	
- Statement Rega	rding Fed sponsored R&D	C.	Г	Stateme	ent verifying is	dentity of ab	ove copies	
- Reference to Mi	crofiche Appendix							
 Background of t 	he Invention			ACCOM	PANYING A	PPLICATIO	N PARTS	
- Brief Summary o	the Invention	7.	П	Assignme	nt Papers (co	ver sheet &	document(s))	
 Brief Description Detailed Descript 	of the Drawings (if filed)	8.	П		§3.73(b) State is an assigned		Power of Attorney	
- Claim(s)		9.	П	English Tr	anslation Do	cument (if a	oplicable)	
- Abstract of the Di	sclosure	10.	П	Information Statement	Disclosure IDS)/PTO-144	9	Copies of IDS Cital	tions
3. Drawing(s) (35 U.S	S.C. 113) [Total Sheets]	11.	П	Preliminar	y Amendmer	nt	_	
Oath or Declaration	[Total Pages 3]	12.	x	(Should be s)	ceipt Postcar pecifically itemize	d)		
a. X Unexecuted	(original or copy)	13.		*Small Entrly Statement(s) (PTO/SB/09-	12)	Status still prop	in prior application, or and desired	
b. Copy from a p	rior application (37 C F R. § 1.63(d)) divisional with Box 16 completed)	14.	П	Certified Co	opy of Priority D	Occument(s) ed)		
	ETION OF INVENTOR(S) ned statement attached deleting	15.		Other.				
inve	entor(s) named in the prior application, see							
NOTE FOR ITEMS 1 & 13: IN ORD	C.F.R. §§ 1 63(d)(2) and 1.33(b) ER TO BE ENTITLED TO PAY SMALL ENTITY ENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT			-		10.0		
IF ONE FILED IN A PRIOR APPLIC	ATION IS RELIED UPON (37 C.F.R. § 1.28).	_						
16. If a CONTINUING APPL	ICATION, check appropriate box, and supply the	e requisi	te ınfo.	rmation below :	and in a prelimina	ry amendment:		
Continuation Di	visional Continuation-in-part (CIP)		of		ion No:			
Prior application informat					/ Art Unit:	eath as doalo	ention is supplied up	ndor
Box 4b, is considered a part of	NAL APPS only: The entire disclosure of the disclosure of the accompanying cont	inuatio	n or	divisional ap	plication and	is hereby inc	corporated by refere	nce.
The incorporation can only be	relied upon when a portion has been inad	verten	ly on	nitted from t	he submitted	application p	arts.	_
	17. CORRESPO	NDEN	CE A	DDRESS				
☑ Customer Number or Ba	r Code Label (Insert Customer No. or.		r code	s label here)	or 🗆	Customer a	ddress below	
Name		.00	J,					
TVUITIO .	PATENT	TRADEHA	RK OF	FICE				
Address								_
City	S	State	T			Zip Code		
Country	Teleph	one	16	510-278-4	1929	Fax	610-278-6548	,
Name (Print/Type)	Steven J. Trzaska	Rec	istra	tion No. (Att	omev/Agent)	36.296		1
Signature	Citoron o. Hizadada	1			Date		per 19, 2000	1

Burden Hour Statement: this form is estimated to take 0.3 bears to complete. Time will vary depending upon the needs of the individual case comments on the amount of time you are required to principle this form should be sent to the Chief information Officer, Patent and Trademise Office, Washington, DC 20231. DO NOT SEND FEES OF COMPLETED PORMS TO THIS ADDRESS. SEND TO Assistant Commissioner for

Patents, Box Patent Application, Washington, DC 2023

FEE TRANSMITTAL	Sports to a contection of miles	Complete if Known	T
	Application Number		2
for FY 2000	Filing Date	w.Z	2 =
Patent fees are subject to annual revision. Small Entity payments must be supported by a small entity statement.	First Named Inventor	Anderson, Kevin W.	
otherwise large entity fees must be paid. See Forms PTO/SB/09-12. See 37 C.F.R. §§ 1.27 and 1.28.	Examiner Name	265	2
TOTAL AMOUNT OF PAYMENT (\$)	Group/Art Unit	Ť	F
834.00	Attorney Docket No.	M 6560 OS/OAPT	

834.00		Attori	ney Do	ket No.	M 6560 OS/OAPT	
METHOD OF PAYMENT (check one)				FEE	CALCULATION (continued)	
The Commissioner is hereby authorized to charge indicated fees and credit any over payments to:	3. 4	DDIT	IONA	L FEE	S	
Deposit Account Number 50-1177 Order No. <u>00-0548</u>	Large Fee Code 105	Entity Fee (\$) 130	Small Fee Code 205	Entity Fee (\$)	Fee Description F Surcharge – late filing fee or oath	Fee Paid
Deposit O	1 127	50	205	65 25	Surcharge – late provisional filing fee or cover sheet.	
Account Cognis Corporation	139	130	139	130	Non-English specification	
Charge Any Additional Fee Required Under 37 CFR §§ 1.15 and 1.17	147	2,520	147	2,520	For filing a request for reexamination	
2. Payment Enclosed:	112	920*	112	920*	Requesting publication of SIR prior to Examiner action	
☐ Check ☐ Money Order ☐ Other	113	1,840*	113	1,840*	Requesting publication of SIR after Examiner action	
FEE CALCULATION	115	110	215	55	Extension for reply within first month	
BASIC FILING FEE	116	380	216	190	Extension for reply within second month	
Large Entity Small Entity Fee Fee Fee Fee Fee Description Fee Paid	117	870	217	435	Extension for reply within third month	
Gode (S) Code (S)	118	1,360	218	680	Extension for reply within fourth month	
101 690 201 345 Utility filing fee \$690.0	0 128	1,850	228	925	Extension for reply within fifth month	
106 310 206 155 Design filing fee	119	300	219	150	Notice of Appeal	
107 480 207 240 Plant filing fee	120	300	220	150	Filing a brief in support of an appeal	
108 690 208 345 Reissue filing fee	121	260	221	130	Request for oral hearing	
114 150 214 75 Provisional filing fee	138	1,510	138	1,510	Petition to institute a public use proceeding	
N	140	110	240	55	Petrition to revive – unavoidable	
SUBTOTAL (1) (\$) 690.00	141	1,210	241	605	Petrition to revive - unintentional	
	142	1,210	242	605	Utility issue fee (or reissue)	
2. EXTRA CLAIM FEES	143	430	243	215	Design issue fee	
Extra Claims Fee from below Fee Paid	144	580	244	290	Plant issue fee	
Total Claims 28 -20**= 8 X \$18.00 = 144.00	122	130	122	130	Petitions to the Commissioner	
ndependent 3 - 3**= 0 X \$78 00 = 0.00	123	50	123	50	Petitions related to provisional applications	
Multiple Dependent \$260.00 =	126	240	126	240	Submission of Information Disclosure Stmt	
"or number previously paid, if greater, For Reissues, see below	581	40	581	40	Recording each patent assignment per property (times number of properties)	
Large Entity Small Entity	146	690	246	345	Filing a submission after final rejection	
Fee Fee Fee Fee Fee Description Code (\$) Code (\$)	149	690	249	345	(37 CFR § 1.129(a)) For each additional invention to be examined (37 CFR § 1.129(b))	
103 18 203 9 Claims in excess of 20						
102 78 202 39 Independent claims in excess of 3	Other	fe <i>e</i> (spe	ecify)			
104 260 204 130 Multiple dependent claim, if not paid 109 78 209 39 "Reissue independent claims over	011					
110 18 210 9 "Ressue dams in excess of 20	Other	fee (spe	ecify) _			
and over original patent						
SUBTOTAL (2) (\$) 144.00	*Re	duced by	Basic F	lng Fee F	Paid SUBTOTAL (3) (\$) 0	0.00

SUBMITTED BY				Compl	ete (if applicable)
Name (Print/Type)	Steven J. Trzaska	Registration No. (Attorney/Agent)	36,296	Telephone	610-278-4929
Signature				Date	September 19, 2000
W	ARNING:				

WARNING
Information on this form may become public. Credit card information should not be
included on this form. Provide grelit card information and authorization on Pro-2038.

Burden Hour Stelement: This time is estimated to take of 25 host to complete. Time will vary depending upon the needs of the individual case, Any comments on the amount of
time you are mounted to complete his form should be sent to the Chief Information Officer, Patient and Trademark Officer, Washington, DC 20231. DO NOT SEND FEES OR
COMPLETED FORMS TO THIS ACROPIES. SORTHOT O Asserts Commissioner for Pleasing Washington, DC 20231.

15

20

25

PATENT Case M 6560 OS /OAPT

IMPROVED FERMENTATION PROCESS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of copending provisional application serial number 60/156,791 filed on September 30, 1999.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was funded, at least in part, under a grant from the Department of Commerce, NIST-ATP Cooperative Agreement Number 70NANB8H4033. The Government may therefore have certain rights in the invention.

BACKGROUND OF THE INVENTION

This invention relates to an improved fermentation medium and process for making an aliphatic polycarboxylic acid using said medium.

Long-chain alpha, omega-dicarboxylic acids, i.e., those having a carbon number of 9 or higher, are used as raw materials in the synthesis of a variety of chemical products and polymers.

Diacids with carbon numbers greater than four are currently produced almost exclusively by nonbiological conversion processes. These types of chemical processes for the production of diacids have a number of limitations and disadvantages. Each process is restricted to the production of diacids of specific carbon chain lengths, based on the starting material used. For example, the dodecandioic acid process begins with butadiene, therefore the products of this reaction process are limited to acids with chain lengths in multiples of four. In addition, the processes are based on nonrenewable petrochemical feedstocks, and the multireaction conversion process produces

25

30

35

unwanted byproducts which result in yield losses, heavy metal wastes, and nitrogen oxides which must be destroyed in a reduction furnace.

Biological conversion processes for the production of diacids have a number of potential advantages relative to the existing non-biological conversion processes. Primary among these is the use of renewable feedstocks as starting materials and the ability to produce the diacid without the generation of hazardous chemical byproducts which necessitate costly waste disposal processes.

Another important advantage achieved by using a biological process is that such a process can easily be adapted to produce a wide variety of diacids using the same biocatalyst and the same equipment. Because current organic chemical syntheses are suited to the production of only a single diacid, the sythesis of several different diacids would require the development of a new synthetic scheme for each diacid. On the other hand, a yeast biocatalyst can be used to produce diacids of varying lengths using the same equipment, media and protocols merely by providing a different substrate to the yeast.

U.S. Patent No. 6,004,784, the entire contents of which is hereby incorporated by reference, discloses a semi-synthetic fermentation medium which employs corn steep liquor and brewers yeast extract in order to reduce the cost of conventional fermentation media which contain expensive, highly standardized yeast extracts and yeast nitrogen bases.

The problem associated with the use of such inexpensive substitutes is many fold. They result in a fermentation broth having significant odor emissions when sparged with air. Particulate matter, especially combined with high levels of bacteria, contained in corn steep liquor and crude yeast extracts make them difficult to sterilize and contribute to the bioburden on media sterilization equipment. These substitutes also contain many unmetabolizable components that contribute to color and color stability problems which need to be attended to using additional purification steps with their incumbent product losses. The selection of these substitutes, while lowering media cost, add additional process costs. Consequently, there remains a need for a low-cost, biofermentation medium which provides nutrients to support growth of the yeast biocatalysts permitting high specific productivity of polycarboxylic acids, polyols, and polyhydroxy acids.

SUMMARY OF THE INVENTION

15

20

25

30

35

The present invention is directed to a fermentation medium and process for making polycarboxylic acids, polyols, and polyhydroxy acids using said medium. The fermentation medium contains:

- (a) a source of metabolizable carbon and energy:
- (b) a source of inorganic nitrogen;
- (c) a source of phosphate:
- (d) at least one metal selected from the group consisting of alkali metals, alkaline earth metals, transition metals, and mixtures thereof; and
- (e) a source of biotin, substantially free of particulate matter and $$\operatorname{\textsc{10}}$$ bacteria.

The present invention is also directed to a process for making polycarboxylic acids, polyols and polyhydroxy acids involving:

- (a) providing an organism capable of producing a polycarboxylic acid, a polyol or a polyhydroxy acid;
 - (b) providing a substrate capable of being converted into a polycarboxylic acid, a polyol or a polyhydroxy acid by the organism;
 - (c) providing a fermentation medium containing:
 - (i) a source of metabolizable carbon and energy;
 - (ii) a source of inorganic nitrogen:
 - (iii) a source of phosphate;
 - (iv) at least one metal selected from the group consisting of alkali metals, alkaline earth metals, transition metals, and mixtures thereof; and
- (v) a source of biotin, substantially free of particulate matter and bacteria; and
 - (d) fermenting the organism in the fermentation medium.

DESCRIPTION OF THE INVENTION

All numbers expressing quantities of ingredients and/or reaction conditions are to be understood as being modified in all instances by the term "about".

Through the judicious selection of alternative nutrients that support microbial growth, the invention provides a medium which allows important polyfunctional materials to be produced commercially, in large quantity, using

15

20

25

30

35

a biological conversion process. The invention provides a low cost alternative to prior art media and methods for its use that is easily prepared and sterilized, has a low degree of odor, provides a lower contribution of impurities loading in subsequent downstream processes, yet results in the production of product in at least as high a yield as conventional prior art methods. It is surprising that the myriad of undefined components comprising corn steep liquor and yeast extract can be replaced by relatively few components without sacrificing the quality of microbial growth and productivity of polyfunctional compounds.

Thus, according to one aspect of the present invention, there is provided an economical fermentation medium capable of facilitating the bioconversion of various types of organic substrates. The fermentation medium contains the following necessary components: (i) a source of metabolizable carbon and energy; (ii) a source of inorganic nitrogen; (iii) a source of phosphate; (iv) at least one metal selected from the group consisting of alkali metals, alkaline earth metals, and mixtures thereof, and (v) a source of biotin, substantially free of particulate matter and bacteria. These materials alone satisfy the basic nutritional requirements for growing the microorganism yet produce biomass of sufficient quantity and quality to conduct the bioconversion

Suitable sources of metabolizable carbon and energy include but are not limited to glucose, fructose, maltose, glycerol, sodium acetate, methanol, short chain alcohols, and mixtures thereof. The preferred source of metabolizable carbon and energy is glucose, preferably a liquid glucose syrup, for example, 95% dextrose- equivalent syrup, or even lower dextrose-equivalent syrups. Such materials contain small amounts of disaccharides, trisaccharides, and polysaccharides which can be hydrolyzed during the fermentation by the addition of an amylase enzyme such as amylase, glucoamylase and cellulase. Thus, glucose can be provided *in situ* in a reaction simultaneous with the biooxidation.

Inorganic sources of nitrogen include but are not limited to alkali metal nitrates such as sodium or potassium nitrate, ammonium salts such as ammonium sulfate, ammonium chloride, ammonium nitrate and ammonium acetate. A prefered inorganic nitrogen source is ammonia or ammonium hydroxide.

Another necessary component of the fermentation medium of the present invention is a source of phosphate which includes any phosphate-containing compounds. Examples thereof include, but are not limited to,

15

20

25

30

35

potassium phosphate, sodium phosphate, and ammonium phosphate. A particularly preferred source of phosphate for use in the present invention is potassium phosphate.

Suitable metals for use in the fermentation medium include alkali metals, alkaline earth metals, transition metals, and mixtures thereof. Particularly preferred metals include potassium, calcium, and magnesium, in combination.

The last critical component of the fermentation medium is biotin which is substantially free of particulate matter and bacteria.

Any of the medium components may be added as a part of the initial sterile media charge, sterilized as concentrated aqueous solutions for later addition to the fermentation medium, or may be included in large excess in the fermentor inoculum for carry-over into the production fermentor.

In order to avoid problems associated with odor emission, color instability, and contamination, it is imperative that the biotin be free of particulate matter and bacteria which promote such problems and require additional process purification steps. The quantity of biotin required will be several orders of magnitude smaller than the organic nitrogen sources it replaces in prior art media.

The fermentation medium of the present invention contains in aqueous solution: (a) from about 10 g/l to about 60g/l, preferably from about 20 g/l to about 40 g/l of a source of metabolizable carbon and energy, preferably glucose; (b) from about 50 ppm to about 2000 ppm, preferably from about 50 ppm to about 250 ppm, and most preferably from about 100 ppm to about 250 ppm, of nitrogen provided by the inclusion of an inorganic nitrogen source; (c) from about 1g/l to about 10 g/l, preferably from about 1 g/l to about 7 g/l, of a source of phosphate, preferably potassium phosphate; (d) from about 0.01 g/l to about 2 g/l, preferably from about 0.01 g/l to about 1 g/l of at least one metal selected from the group consisting of alkali metals, alkaline earth metals, and mixtures thereof, preferably a mixture of calcium and magnesium; and (e) from about 1 $\mu g/l$ to about 2000 $\mu g/l$, preferably from about 4 $\mu g/l$ to about 200 $\mu g/l$, of blotin, wherein the biotin is substantially free of deleterious particulate matter and bacteria.

The water present in the fermentation medium may be a process water purified by distillation, deionization, or softening. Preferred sources of water include those from a municipal water distribution system, a process recycle stream, or well water wherein adjustments in mineral content may need to be

15

20

25

30

35

taken into account for minerals already contained in these sources of water. For example, the water together with other required ingredients may already contain sufficient mineral components to provide all or substantially all the required minerals for growth of the organism.

The fermentation media of the present invention satisfies the basic nutrient requirements for the growth of the microorganism, while minimizing the addition of organic and inorganic components that require separation from the product for disposal as waste and contribute to process odor emissions. In contrast to fermentation media compositions known in the art, the media of the present invention contains no particulate matter, is particularly amenable to continuous sterilization in an automated medium preparation process and is inexpensive. Despite the nutritionally lean media compositions provided by the present invention, there is broad flexibility in formulating the media while achieving high productivities of polycarboxylic acids, polyols, and polyhydroxy acids.

Various types of auxiliary components may also be employed in the fermentation medium of the present invention in order to further enhance the biofermentation process. Examples thereof include, but are not limited to, various types of trace metals, chelating agents, anti-foaming agents, and the like.

The fermentation media of the present invention may be used with any polycarboxylic acid-producing, polyol-producing and polyhydroxy acid-producing yeast, and a wide variety of substrates. For example, in the event that the desired product is a polycarboxylic acid, such as a diacid, any type of fatty acid, fatty acid ester, or alkane substrate may be used. Examples of suitable substrates for the production of diacids include, but are not limited to, lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, palmitoleic acid and methyl esters thereof, and mixtures thereof. Examples of alkanes include dodecane, dodecene, tridecane, tetradeance, octadecane, and the like.

Thus, according to another aspect of the present invention, there is provided a process for making a polycarboxylic acid, a polyol, or a polyhydroxy acid. For examplary purposes only, the process of the present invention will be described with reference to making polycarboxylic acids, and specifically diacids, as the desired end product.

The process may be operated over any pH range where the microorganism can grow and catalyze the desired conversion reaction. A prefered and especially advantageous pH range is in the acidic regime, i.e., a pH of about 7 or less. While the low pH improvement can be applied to any

15

20

25

35

fermentation process, it is especially advantageous when applied to a fermentation process which produces polycarboxylic acids. Suitable pH control reagents are ammonia, ammonium hydroxide solution, concentrated potassium or sodium hydroxide.

The organic substrate can be any organic compound that is biooxidizable to a mono- or polycarboxylic acid. Such a compound can be any saturated or unsaturated aliphatic compound or any carbocyclic or heterocyclic aromatic compound having at least one terminal methyl group, a terminal carboxyl group and/or a terminal functional group which is oxidizable to a carboxyl group by biooxidation. A terminal functional group which is a derivative of a carboxyl group may be present in the substrate molecule and may be converted to a carboxyl group by a reaction other than biooxidation. For example, a lipase enzyme, for example, can be added during the fermentation step to liberate free fatty acids from an otherwise non-metabolizable ester.

Alkanes are a type of saturated organic substrate which are useful in practicing the process according to the invention. The alkanes can be linear or cyclic, branched or straight chain, substituted or unsubstituted. Particularly preferred alkanes are those having from about 4 to about 25 carbon atoms, examples of which include but are not limited to butane, hexane, octane, nonane, dodecane, tridecane, tetradecane, octadecane and the like.

Examples of unsaturated organic substrates which can be used in the process according to the invention include but are not limited to internal olefins such as 2-pentene, 2-hexene, 3-hexene, 9-octadecene and the like; unsaturated carboxylic acids such as 2-hexenoic acid and esters thereof, oleic acid and esters thereof including a triglyceryl esters having a relatively high oleic acid content, erucic acid and esters thereof including triglyceryl esters having a relatively high erucic acid content, ricinoleic acid and esters thereof including triglyceryl esters having a relatively high ricinoleic acid content, linoleic acid and esters thereof including triglyceryl esters having a relatively high oleic acid content; unsaturated alcohols such as 3-hexen-1-ol, 9-octadecen-1-ol and the like; unsaturated aldehydes such as 3-hexen-1-al, 9-octadecen-1-al and the like. In addition to the above, the organic substrate which can be used in the process according to the invention include alicyclic compounds having at least one internal carbon-carbon double bond and at least one terminal methyl group, a terminal carboxyl group and/or a terminal functional group which is oxidizable to a carboxyl group by biooxidation. Examples of such compounds include but are not limited to 3,6-dimethyl-1,4-

20

25

30

35

cyclohexadiene; 3-methylcyclohexene; 3-methyl-1, 4-cyclohexadiene and the like.

Examples of the aromatic compounds that can be used in the process according to the invention include but are not limited to arenes such as o-, m-, p-xylene; o-, m-, p-methyl benzoic acid; dimethyl pyridine, and the like. The organic substrate can also contain other functional groups that are biooxidizable to carboxyl groups such as an aldehyde or alcohol group. The organic substrate can also contain other functional groups that are not biooxidizable to carboxyl groups and do not interfere with the biooxidation such as halogens, ethers, and the like.

A cosubstrate may be provided to the culture during that stage of the fermentation when the culture is actively oxidizing the substrate to products. This is especially true if no net useable carbon or energy can be recovered from the substrate reaction itself, then this carbon and energy is provided by the cosubstrate.

The cosubstrate is a fermentable carbohydrate such as glucose, fructose, or maltose; or other fermentable organic compound, for example, glycerol, sodium acetate, methanol, or short chain alcohols; or mixtures thereof. The preferred cosubstrate is glucose, preferably a liquid glucose syrup, for example, 95% dextrose- equivalent syrup, or even lower dextrose-equivalent syrups. Such materials contain small amounts of disaccharides, trisaccharides, and polysaccharides which can be hydrolyzed during the fermentation by the addition of an amylase enzyme such as α -amylase, glucoamylase and cellulase. Thus glucose can be provided $in\ situ$ in a reaction simultaneous with the biooxidation.

It is convienient, but not absolutely necessary, if the carbon and energy source used to grow the biomass is the same as the cosubstrate used to drive the oxidative conversion reaction. The practical benefit is that fewer raw materials need be handled and the various stages of the fermentation can be well integrated. Thus a single cosubstrate sterilization and delivery system can be used to deliver both the carbon and energy source to grow the biomass and the cosubstrate to drive the oxidation reaction.

The microorganism that can be used in the process according to the invention is any microorganism capable of biooxidizing the substrate as defined herein. The microorganism can be any microorganism in which beta oxidation is partially or completely blocked by disruption, inactivation or deletion of one or more acyl CoA oxidase gene(s); any yeast in which beta oxidation is partially or completely disrupted by inactivation or deletion of one

15

20

25

30

or more acyl CoA oxidase gene(s); any Candida strain where beta oxidation is partially or completely disrupted by inactivation or deletion of one or more acyl CoA oxidase gene(s). When the fermentation process involves the biooxidation of a substrate to a carboxylic acid, the microorganism will normally be a yeast. Such a microorganism is capable of oxidizing the substrate in such a way that the mono- and/or dicarboxylic acids formed are not further oxidized by degradation leading to chain shortening. However, the process according to the invention pertains to the use of any microorganism.

Yeast strains known to excrete alpha, omega-dicarboxylic acids as a by-product when cultured on alkanes or fatty acids as the carbon source are set forth in U.S. patent 5,254,466, the entire contents of which are herein incorporated by reference. These strains are partially or completely oxidation-blocked strains; that is, they are genetically modified so that the chromosomal POX4A, POX4B and both POX5 genes have been disrupted. The substrate flow in this strain is redirected to the omega-oxidation pathway as the result of functional inactivation of the competing \$\mathbb{G}\$-oxidation pathway by POX gene disruption. A completely oxidation-blocked strain is a \$\mathbb{C}\$. tropicalis strain, strain H5343 (ATCC 20962), which is described in U.S. patent 5,254,466.

Another suitable strain is one in which one or more reductase genes are amplified resulting in an increase in the amount of rate-limiting omegahydroxylase through P450 gene amplification and an increase in the rate of substrate flow through the ω -oxidation pathway. Strains which selectively increase the amount of enzymes known to be important to the oxidation of fatty acids are also preferred. Such strains contain increased copies of the CYP and CPR genes. These genes have been identified as those relating to the production of the ω -hydroxylase complex catalyzing the first step in the oxidation pathway. Strain HDC1 is an example of a strain that contains multiple copies of the CYP 52A2A gene integrated into the genome of strain H5343. This strain and similar strains are described in copending application serial number 60/083,798, filed on 05/01/98, the entire contents of which are incorporated herein by reference. Other strains that can be used with the methods of this invention are *C. tropicalis* strains HDC1, HDC5, HDC10, HDC15, HDC20, HDC23, HDC 23-1, HDC 23-2, and HDC 23-3 as are

15

20

25

30

described in PCT/US99/20797, the entire content of which is hereby incorporated by reference.

Suitable sources of inorganic nitrogen include any ammonium-containing compounds. Examples thereof include, but are not limited to, alkali metal nitrates or ammonium salts such as ammonium sulfate, ammonium chloride, ammonium nitrate and ammonium acetate. Prefered nitrogen sources are ammonium salts or compounds that generate ammonia through the metabolic action of the organism like urea. A particularly preferred source of inorganic nitrogen for use in the present invention is ammonia.

The fermentation process can be modified by utilizing a triglyceride fat or oil as the source of both the organic substrate and cosubstrate. A lipase, formulated with the fermentation broth, hydrolyzes or splits the fat or oil into fatty acids and glycerine. Glycerine consumption by the organism serves to drive the splitting reaction to completion while supplying the energy necessary to convert the free fatty acids to their corresponding dibasic acids. Lipases that are oleo-specific are particularly preferred. Oleo-specific lipases exhibit a high selectivity for a triglyceride having a high oleic acid content and selectively catalyze the hydrolysis of the oleate ester groups. Examples of such oleo-specific lipases include but are not limited to the lipases produced by Pseudomonas sp, Humicola lanuginosa, Candida rugosa, Geotrichum candidum, and Pseudomonas (Burkholderia). A particularly preferred lipase is UNLipase from Geotrichum candidum ATCC No. 74170 described in U.S. patent 5,470,741, the entire contents of which are incorporated herein by reference.

The fermentation step is preferably carried out in three stages, growth phase, induction phase, and conversion phase. Each phase may be operated at the same or different fermentor conditions of temperature, pH, aeration, etc. Growth phase begins when the cell culture is introduced into the fermentor and a rapid phase of growth ensues. Growth may be exponential or sub-exponential depending on the composition of the medium employed. This continues until the culture reaches linear growth as measured by a decrease in oxygen consumption. Linear growth occurs when

growth is limited by the rate of addition of a key nutrient to the medium; thus growth becomes proportional to the rate at which the limiting nutrient is supplied. Usually, in the process of this invention, the key nutrient will be the cosubstrate. The second phase is the induction phase. In the induction phase key metabolic activities are initiated that begin the desired conversion of substrate to product. The culture can be maintained in a linear growth phase for a period of time before inducing the culture. The inducing agent will usually be the substrate itself, but for compunds that do not initiate their own conversion, another inducing agent may be used in concert with the substrate. The induction phase can be used to transition the rapid growth phase to the conversion phase. The fermentation moves into the next phase, conversion phase, when the substrate is converted to product.

During the conversion phase, the fermentation broth is in an acidic pH range of between 2 and 7. The preferred operating pH range is from about 3.5 to 7.0, even more preferred range is from about 5.0 to about 6.5. pH may be controlled by automatic titration using a strong base, examples of which are, sodium hydroxide solution, potassium hydroxide solution, ammonium hydroxide solution, or ammonia gas. Note that by operating the fermentation in this pH regime compared to prior art methods that ammonia can be used for pH control since it will react to form the non volitile aqueous ammonium ion. Prior art methods for making dicarboxylic acid when operated in the alkaline pH regime would not effectively make the aqueous ammonium ion thus causing undesirable emissions of ammonia vapor in the offgas and toxic ammonia accumulation in the broth. The fermentation can be carried out at a temperature of from about 26 °C to about 40 °C.

In the first stage, a culture medium is inoculated with an active culture of beta-oxidation blocked microorganism such as a beta-oxidation blocked $\underline{\mathbb{C}}$. tropicalis strain where a period of rapid exponential growth occurs. The pH of the culture medium is controlled by the addition of base, examples include but are not limited to ammonium hydroxide, potassium hydroxide, sodium hydroxide or ammonia gas. The cosubstrate addition to the fermentor is preferrably a fed-batch addition during the conversion phase. The end of exponential growth phase is marked by a depletion of glucose, a rapid increase in dissolved oxygen, and, most sensitively, by a rapid increase in offgas oxygen and decrease in offgas CO₂. In the absence of an inducing substrate, biomass will continue to accumulate at a rate proportional to the glucose feed rate (i.e., linear growth). It is desirable to initiate the conversion phase either at the end of exponential growth or at a point when the desired

15

20

25

35

biomass level has been attained. If the oxygen transfer characteristics of the production vessel are such that the desired biomass level can not be reached by maintaining the culture in exponential growth, then a combination of exponential growth followed by a linear glucose-limited growth phase (induction phase) can be used to achieve a given biomass level.

To maintain the culture in a healthy state during the transition to glucose limited linear growth, it may be desirable to start a glucose feed during growth so that glucose will always be available to the culture. The glucose level in the initial medium charge is selected based on the total glucose desired for growing the microorganism less the amount that will be fed during growth. The glucose feed may be started at any time from the beginning of growth. However because there is some variation in inoculum quality and lag phase after inoculating the production vessel, it is preferred to start the glucose feed during exponential growth phase at some predetermined biomass concentration, as judged by direct biomass measurements, or indirect methods like optical density, carbon dioxide evolution rate, or oxygen uptake rate. The glucose may be a corn syrup refined or unrefined. The glucose may be fed from the top of the vessel through the vapor space as a continuous stream or as a continuous series of pulses or impulses. The glucose feed may also be fed into the vessel under the surface especially into a region of high shear near the agitator to attain a rapid distribution throughout the vessel. In large vessels there may be multiple glucose feed points within the same vessel.

The conversion phase where the substrate is oxidized is initiated by adding an inducer and the substrate containing an oxidizable methyl group. In the case of alkanes, fatty acids, fatty acid methyl esters and fatty acid salts; these substances and combinations thereof induce their own oxidation to dicarboxylic acids and may be useful for inducing the oxidation of other substances. The substrate may be added batchwise or as a continuous feed or as a series of continuous pulses or impulses. The substrate may be added from the top of the vessel through the vapor space or under the surface especially into a region of high shear. In large vessels there may be multiple substrate feed points. The oxidation is conducted at an acidic pH less than 7 and preferable near or below the pKa of the carboxyl group being formed by the oxidation or is present in the substrate.

It is preferable to progressively decrease the glucose feed rate to the fermentor as the conversion proceeds to prevent the accumulation of biomass and triacylglycerol esters. With the conversion of commercial oleic acid, the

20

25

30

35

accumulation of intracellular storage vacuoles is a useful indicator of the glucose feed rate needing adjustment. Adjustments are done by decreasing the feed rate by about 5% to 25% and usually about 10% per each 24 hours of the conversion phase. Other indicators could also be used such as base utilization rate, CO2 evolution rate, or respiratory quotient.

One problem often encountered in the fermentation process is the formation of carboxylic acid soaps and foaming as a result of working in the traditional alkaline pH range. In an alkaline environment, carboxylic acids form soaps which lead to unwanted foaming of the fermentation broth. The result of the soap formation is a decrease in the pH of the broth which is adjusted by adding a base such as caustic to the broth to maintain the pH of the broth at the desired value. Also, addition of glucose to the basic environment results in formation of carbonates that affect the pH of the fermentation broth. To compensate for the effects on the pH of soap and carbonate formation, large quantities of base must be added to maintain the pH of the broth at the desired level.

It has surprisingly been found that by carrying out the fermentation process in an acidic pH range of from 2 to 7, preferably from 3 to 7 and even more preferably for 5 to 6.5, rather than a caustic pH range, foaming is substantially reduced and the formation of carbonate from carbon dioxide produced during the metabolism of glucose is reduced thereby substantially reducing the amount of raw material needed to control the pH of the fermentation broth. By carrying out the fermentation process at an acidic pH. these problems are substantially reduced resulting in decreased amounts of base used during the fermentation process. There is a net consumption of cation equivalents over anion equivalents during the growth reaction that contributes to a depression of culture medium pH. It is desirable to control the medium pH within the range from 2-7, and preferably from about 3-6.5 during growth by the addition of base on pH control. Among the bases useful for this purpose are ammonium hydroxide, ammonia, sodium hydroxide, and potassium hydroxide. It is desirable to select a base composition that provides one or more of the major nutrients consumed during growth such as ammonium hydroxide, ammonia, or potassium hydroxide. The growth medium formulation is adjusted to take into account the addition of the pH control reagent. Using NH₄OH or ammonia gas to control the pH reduces the number of raw materials needed to grow the microorganisms by combining the pH control agent and the nitrogen source into one raw material added to the fermentor.

With respect to using nitrogen sources such as ammonia or ammonium hydroxide it is necessary to have only 250 ppm or less nitrogen source present in the medium to initiate growth and the consumption of these pH control reagents. The ammonia concentration will therefore remain nearly constant during growth of the culture. Thus the desired concentration of ammonia in the medium at the time of induction can be conveniently preselected by adding ammonia or ammonium salts to the initial medium charge. Useful sources of ammoniacal nitrogen for the initial fermentor charge include ammonia phosphate, ammonium sulfate, ammonium nitrate, ammonia, urea, and ammonium hydroxide.

Another aspect of the invention relates to a formulation of a fermentation medium useful for propagating Candida tropicalis and gives a high productivity for converting substrates having oxidizable methyl groups to carboxylic acids. By adjusting feed rates of materials into the fermentation broth, the growth of the microorganism can be controlled. The major nutrients consumed during growth of the microorganism on glucose are ammoniacal nitrogen, potassium, magnesium, phosphate, and sulfate. Sodium and calcium are not consumed, however calcium should be present at a concentration of about 5-50 ppm or more to obtain normal growth depending on the formulation of the inoculum medium. Trace minerals and biotin are also included in the medium. The biotin may be a relatively pure grade or supplied as a more crude grade such as in biotin yeast, yeast extracts, or corn steep liquor.

Useful sources of phosphate are ammonium phosphate, potassium phosphate mono, di, and tribasic, sodium phosphate, mono, di, and tribasic, and phosphoric acid. Useful sources of potassium are potassium sulfate, potassium phosphate, mono, di, and tribasic, and caustic potash.

A fermentation aid can be used in the oxidation of alkanes and/or the oxidation of fatty acids. The preferred fermentation aid is a fatty acid ester, a particularly preferred fermentation aid is a methyl ester. One major benefit of the use of such a fermentation control agent is in foam control and controlling the fluid characteristics of the fermentation broth. If product chain length distribution is important in ultimate product performance, it is desirable to select a chain length of the methyl ester that resembles the alkane or fatty acid since the methyl ester will also be converted to product. The ester may be added batch wise or blended into the feed. If blended into the feed, it will comprise about 10% or less and preferably about 1% or less of the feed. Some commercial fatty acids naturally contain some methyl esters and may

20

25

30

35

be used directly as substrates for dicarboxylic acid production. For example, EMERSOL® 267 (Cognis Corporation) is a commercial oleic acid containing about 1% or less methyl esters and has been found to be a good substrate in this invention. A typical technical grade of oleic acid, EMERSOL® 267, that is used in the process has the approximate composition 0.3% C12, 2.4% C14, 0.6% C14:1, 4.7% C16, 4.6% C16:1, 0.2% C17, 0.8% C18, 69.9% C18:1, 10.5% C18:2, 0.3% C18:3.

In the case of alkane oxidations to dicarboxylic acids, it has further been found useful to use a fatty acid or fatty acid salt as a fermentation aid. This results in a better distribution of the alkane throughout the fermentation broth. The fatty acid may be added batchwise or may be formulated into the alkane feed. If formulated into the feed, fatty acid or fatty acid salt will typically comprise about 10% or less and preferably 5% or less of the feed. Here too, if product chain length distribution is important in ultimate product performance, it is desirable to select a chain length distribution of the fatty acid that resembles the alkane since the fatty acid will also be converted to product.

It has surprisingly been found that by maintaining dissolved oxygen concentration levels below about 25% and preferably below 20%, relative to saturation with air, that there is a reduction in the amount of cosubstrate needed during the conversion phase. When the concentration of dissolved oxygen is below these levels, glucose is more efficiently utilized in providing energy for oxidation. Consequences of adding too much glucose include the further accumulation of biomass and triacylglycerol esters with diminished carboxylic acid production during the conversion phase.

Triacylglycerol ester formation, in oleoagenous yeasts, can be controlled or minimized in part by adjustment of the magnesium concentration or ratio of magnesium to phosphate in the initial medium charge. The preferred magnesium concentration is about 0.1 to about 1.0 gm/L and more preferably about 0.2 to about 0.8 g/L concentration in the fermentation incoulum. The preferred phosphate:magnesium ratio is 20:1 to about 2:1, preferably 15:1 to about 3:1.

It has also surprisingly been found that the fermentation broth during the conversion phase accumulates a heat-labile catalase-like activity that consumes H2O2. This may interfere with glucose concentration assays that measure H2O2 produced from glucose oxidase.

A preferred embodiment of the process according to the invention is the preparation of 9-octadecenedioic acid by biooxidation of oleic acid. While

any grade of oleic acid can be used as the substrate, a typical technical grade oleic acid consists of the following carboxylic acids: 0.42% C₁₂; 2.7% C₁₄; 0.86% C₁₄:1 6.3% C₁₆; 4.6% C₁₆:1; 0.93% C₁₇; 2.8 C₁₈; 71.8% C_{18:1}; 8.3% C_{18:2}; 0.58% C_{18:3}. The oleic acid can also be a high oleic acid grade obtained from a fatty oil of a Helianthus annuus (sunflower seed oil) species described, for example, in U.S. Pat. No. 4,627,192, the entire contents of which are incorporated herein by reference. Other high oleic acid varieties of oilseeds may also be used in this process. Such oils are very rich in oleic acid and contain at least 70 -80% by weight of oleic acid.

The present invention will be better understood from the examples which follow, all of which are intended for illustrative purposes only, and are not meant to unduly limit the scope of the invention in any way. In all the examples, component concentrations are shown in their anhydrous forms. Commercially available hydrates of the listed components may be used if the concentration is appropriately adjusted to include waters of hydration.

EXAMPLE 1

Fermentation media in accordance with the present invention were prepared, the components of which are listed in Tables 1 and 2, below.

Table 1

Table 1
Concentration (g/L)
27.0
7.0
5.1
0.5
0.1
0.06
0.023
0.0002
0.0009
0.00007
0.00018
0.00036
0.00072
0.00036
0.00072
Balance
0.8 ml

5 The medium components of Table 1 were heat sterilized in a suitable manner to avoid any precipitation reactions, then combined in a sterile fermentor vessel upon cooling. The complete, uninoculated medium was found to be completely clear with a slight straw color and no strong odors. Addition of the antifoam yielded slight turbidity. Candida tropicalis H5343 - 10 ALK 2-1 was grown under sterile conditions on the medium listed in Table 1 in a stirred, aerated fermentor with an initial liquid volume of 12 L. The sterile culture medium was inoculated with a 5% inoculum of Candida tropicalis H5343 ALK 2-1 and grown at 35°C, pH 5.8 for approximately 10 hours with agitation and aeration rates sufficient to keep the dissolved oxygen above 20%. When the culture stopped growing exponentially and the dissolved 15 oxygen began to rise, the conversion phase was commenced by starting a continuous feed stream of Exxon Developmental Fluid 137 (a hydrocarbon

15

containing approximately 94.4% Tridecane, the balance being primarily dodecane) mixed with the fermentation aids 1.25% Emersol ®267 (a technical grade of oleic acid) and 1.25% Emery® 2203 (a technical grade of methyl tallowate) at a rate of 0.7 g/L/hr. Simultaneously, the temperature in the fermentor was reduced from 35°C to 30°C, the aeration rate was reduced to 0.4 vvm, and 0.4 bar of backpressure was applied to the vessel. The pH was maintained between 5.8 – 5.9 during the growth and conversion phase with 6N KOH. A continuous glucose feed stream was started to the fermentor at a rate of 1.58 g/L/hr glucose when the biomass concentration reached approximately 10g/L. The glucose feed rate during conversion was reduced between 0 – 15 % on a daily basis based upon the microscopic observation and assessment of accumulating storage vacuoles within the yeast cell. 7 ml of PPG (polypropylene glycol) antifoam was added to the fermentor during conversion to control mild foaming. After 50 hours of the conversion phase, the whole broth in the fermentor contained 41.5 g/Kg 1,13-tridecanedioic acid.

EXAMPLE 2

20		Table 2
	Synthetic Production Medium Components	Concentration (g/L)
	Glucose	27.0
	Potassium Phosphate, Monobasic	4.9
	Magnesium Sulfate	0.6
	Calcium Chloride	0.1
	Citric Acid	0.06
	Ferric Chloride	0.023
	Biotin	0.000012
	Trace Metals:	
	Cupric Sulfate	0.00007
	Manganese Sulfate	0.00432
	Zinc Sulfate	0.00072
	Citrate	0.00708
	Water	Balance
	SAG 471 ®Antifoam	0.6 ml

20

25

30

The medium components of Table 2 were sterilized in a suitable manner to avoid any precipitation reactions and combined in a sterile fermentor vessel. The complete, uninoculated medium was found to be completely clear with a slight straw color and no strong odors. Candida tropicalis H5343 HDC 23-3 was grown under sterile conditions on the medium listed in Table 2 in a stirred, aerated fermentor with an initial liquid volume of 12 L. The sterile culture medium was inoculated with a 3% inoculum of Candida tropicalis H5343 HDC 23-3 and grown at 35°C for approximately 12 hours with agitation and aeration rates sufficient to keep the dissolved oxygen above 20%. The pH was adjusted to and maintained between 5.8 - 5.9 during the growth phase with the addition of 6N NH₄OH which was also the source of inorganic nitrogen in the medium. When the culture stopped growing exponentially and the dissolved oxygen began to rise, the conversion phase was commenced by adding an inducing charge and simultaneously starting a continuous feed stream of High Oleic Sunflower Fatty Acids (containing 84.4% oleic acid, 5.2% linoleic acid, 4.7% stearic acid, 3.9% palmitic acid, with the balance comprising small amounts of eicosanoic acid (20:0), eicosaenoic acid (20:1), pentadecanoic acid, lauric acid, and myristic acid) at a rate of 2.0 g/L/hr. Simultaneously, the temperature in the fermentor was reduced from 35°C to 30°C, the aeration rate was reduced to 0.4 vvm, and the pH control reagent was switched from NH₄OH to NaOH. The pH was maintained between 5.8 -5.9 during the conversion phase with 6N NaOH. A continuous glucose feed stream was started to the fermentor at a rate of 1.22 g/L/hr glucose when the biomass concentration reached approximately 10 g/L. The glucose feed rate during conversion was reduced between 0 - 45% on a daily basis based upon the microscopic observation and assessment of accumulating storage vacuoles within the yeast cell. No additional antifoam was used during the conversion phase. After 50 hours of the conversion phase, the whole broth in the fermentor contained 71 g/Kg total dicarboxylic acids.

35

Example 3

Table 3

Synthetic medium formulation	
Medium Components	Concentration (g/L)
O.	
Glucose	20.0
Ammonium Sulfate	0.5
Potassium Phosphate, Monobasic	5.1
Magnesium Sulfate	0.9
NaCl	0.5
Calcium Chloride	0.1
Biotin	0.0002
Trace Metals:	
Boric Acid	0.00075
Cupric Sulfate	0.00006
Potassium lodide	0.00015
Ferric sulfate	0.04
Ferric Chloride	0.0003
Manganese Sulfate	0.0006
Sodium Molybdate	0.0003
Zinc Sulfate	0.0006
Water	Balance
SAG 471®Antifoam	2 drops

The medium components of Table 3 were sterilized in a suitable manner to 5 avoid any precipitation reactions, then combined in a sterile fermentor vessel. The medium was found to be completely clear with a slight color and little odor. Addition of the antifoam yielded slight turbidity. The pH of the medium was initially adjusted to pH 5.8 using 6N ammonium hydroxide solution. Candida tropicalis H5343 (ATCC 20962) was grown on this medium in a 10 stirred and aerated fermentor using a 4% inoculum prepared in a medium having similar composition. Exponential growth began following a brief lag phase. A glucose feed was started at a rate of 1.25 g/L/hr based on the initial medium volume when the culture contained about 5 g/L biomass dry weight as judged by optical density measurements. Exponential growth continued 15 for about nine hours when the glucose became growth limiting with only a slight decrease in exponential growth rate over the period.

15

At the end of this period of rapid growth, the pH control reagent was changed to 6N potassium hydroxide solution and the glucose feed was kept constant for the next 110 hours. During this period, ammonium, which concentration remained constant throughout exponential growth, was depleted from the medium and phosphate concentration dropped to low levels. Biomass continued to accumulate in the medium at a constant linear rate despite the consumption of these key nutrients. Viable cell counts also increased linearly until the ammonium was depleted from the medium, after which they remained constant. Ultimately, the fermentation produced 71.5 g/L biomass dry weight.

Comparative Example 1

Production Medium Components	Concentration (g/L)
Glucose	40.0
Ammonium Sulfate	8.0
Corn Steep Liquor	9.0
Potassium Phosphate, Monobasic	2.0
Potassium Phosphate, Dibasic	1.0
Magnesium Sulfate	0.5
NaCl	0.5
Calcium Chloride	0.1
Trace Metals:	
Boric Acid	0.00075
Cupric Sulfate	0.00006
Potassium lodide	0.00015
Ferric Chloride	0.0003
Manganese Sulfate	0.0006
Sodium Molybdate	0.0003
Zinc Sulfate	0.0006
SAG 471 ®Antifoam	2 drops

The medium components of Comparative Example 1 were sterilized in a suitable manner to avoid any precipitation reactions and combined in a sterile fermentor vessel. The complete, uninoculated medium was found to be very dark with a strong odor characteristic of com steep liquor. Candida tropicalis H5343 (ATCC 20962) was grown under sterile conditions on the

15

20

25

medium listed in Table 3 in a stirred, aerated fermentor with an initial liquid volume of 10 L. The sterile culture medium was inoculated with a 6% inoculum of Candida tropicalis H5343 (ATCC 20962) and grown at 35°C for approximately 9.5 hours with agitation and aeration rates sufficient to keep the dissolved oxygen above 20%. 5 ml of SAG 471 antifoam was added to the fermentor during growth to control foaming in the culture. The pH was maintained between 5.8 - 5.9 during the growth phase with the addition of 6N NH₄OH. When the culture stopped growing exponentially and the dissolved oxygen began to rise, the conversion phase was commenced by starting a continuous feed stream of Emersol®267 (a commercial grade of Oleic acid containing containing 71.8% oleic acid, 8.3% linoleic acid, 6.3 % palmitic acid, 4.6% palmitoleic acid, 2.8% stearic acid, 2.7% myristic acid, 0.93% C17:0 acid, 0.86% myristoleic acid, 0.58% linolenic acid, and 0.42% lauric acid) at a rate of 2.0 g/L/hr. Simultaneously, the temperature in the fermentor was reduced from 35°C to 30°C, the aeration rate was reduced to 1.2 vvm. and the pH control reagent was switched from NH,OH to KOH. The pH was maintained at or above 5.8 during the conversion phase with 6N KOH. A continuous glucose feed stream was started to the fermentor at a rate of 1.8 g/L/hr glucose at the end of the exponential growth immediately preceeding the start of the conversion phase. This same glucose feed rate was maintained to the fermentor throughout the conversion phase. 17 ml of SAG 471 antifoam was added to the fermentor during the first 3 hours of the conversion phase to control very heavy foaming. After 50 hours of the conversion phase, the whole broth in the fermentor contained 64 g/Kg total dicarboxylic acids.

What is claimed is:

- A fermentation medium comprising:
 - (a) a source of metabolizable carbon and energy;
 - (b) a source of inorganic nitrogen;
 - (c) a source of phosphate:
- (d) at least one metal selected from the group consisting of an alkali metal, an alkaline earth metal, transition metals, and mixtures thereof; and
- (e) a source of biotin, substantially free of particulate matter and bacteria.
- The medium of claim 1 wherein the source of metabolizable carbon and energy is glucose.
- The medium of claim 1 wherein the source of inorganic nitrogen is ammonium sulfate.
- The medium of claim 1 wherein the source of inorganic nitrogen is ammonia.
- The medium of claim 1 wherein the source of inorganic nitrogen is ammonium hydroxide.
- The medium of claim 1 wherein the source of phosphate is potassium phosphate.
- 7. The medium of claim 1 wherein the metal is calcium.
- 8. The medium of claim 1 wherein the metal is magnesium.
- 9. The medium of claim 1 wherein the metal is both calcium and magnesium.
- 10. The medium of claim 1 further comprising an antifoam agent.
- 11. The medium of claim 1 further comprising a chelating agent.
- 12. The medium of claim 1 further comprising at least one trace metal.
- 13. A fermentation medium comprising:
 - (a) glucose:
 - (b) an ammonium salt;
 - (c) a phosphate salt;
 - (d) a potassium salt;
 - (e) magnesium sulfate;

- (f) a calcium salt;
- (g) an ironsalt;
- (h) a chelating agent; and
- (i) a trace metal.
- 14. A process for making a polycarboxylic acid, a polyol, or a polyhydoxy acid comprising:
- (a) providing an organism capable of producing a polycarboxylic acid, a polyol or a polyhydroxy acid:
- (b) providing a substrate capable of being converted into a polycarboxylic acid, a polyol, or a polyhydroxy acid by the organism;
 - (c) providing a fermentation medium containing:
 - (i) a source of metabolizable carbon and energy;
 - (ii) a source of inorganic nitrogen;
 - (iii) a source of phosphate;
- (iv) at least one metal selected from the group consisting of an alkali metal, an alkaline earth metal, a transition metal, and mixtures thereof; and
- (v) a source of biotin, substantially free of particulate matter and bacteria; and
 - (d) fermenting the organism and substrate in the fermentation medium.
- 15. The process of claim 14 wherein the organism is *C. tropicalis*.
- 16. The process of claim 14 wherein the substrate is an alkane having from about 4 to about 25 carbon atoms.
- 17. The process of claim 14 wherein the source of metabolizable carbon and energy is glucose.
- 18. The process of claim 14 wherein the source of inorganic nitrogen is ammonium sulfate.
- The process of claim 14 wherein the source of phosphate is potassium phosphate.
- 20. The process of claim 14 wherein the metal is calcium.
- 21. The process of claim 14 wherein metal is magnesium.

- 22. The process of claim 14 wherein the metal is both calcium and magnesium.
- 23. The process of claim 14 wherein the medium further comprises a chelating agent.
- 24. The process of claim 14 wherein the medium further comprises an antifoam agent.
- 25. The process of claim 14 wherein the medium further comprises at least one trace metal.
- 26. The process of claim 14 wherein step (d) is performed at a pH of up to about 7
- 27. The process of claim 14 further comprising maintaining dissolved oxygen concentration levels below about 25% during step (d).
- 28. The process of claim 14 wherein the source of inorganic nitrogen is selected from the group consisting of ammonia, ammonium hydroxide, and mixtures thereof.

Abstract of the Disclosure

A fermentation medium containing: (a) a source of metabolizable carbon and energy; (b) a source of inorganic nitrogen; (c) a source of phosphate; (d) at least one metal selected from the group consisting of an alkali metal, an alkaline earth metal, a transition metal, and mixtures thereof; and (e) biotin, substantially free of particulate matter and bacteria.

Cognis Corporation Patent Department 2500 Renaissance Blvd., St. 200 Gulph Mills, PA 19406

SJT/mc G:/DATA/TRZASKA/M6560sp.doc

(1) (1) (1) (1)
150
1.64
m
1,4
10
(f)
l,J
99
100
10
j. de
40
100

xpress Mail" mailing label numb Fype a plus sign (+) inside this b		Approve Patent and Trademark	ed for use through Office; U.S. DEF	PTO/SB/01 (6 n: 10/31/98 OMB 0651- ARTMENT OF COMME	0032
0010/PTO Rev. 6/95	U.S. Department of Commerce Patent and Trademark Office	Attorney Docket Number	M 6560 OS/	OAPT	1
DECLARA	ION FOR	First Named Inventor	Anderson, l	Kevin W.	_
UTILITY OF		сом	PLETE IF KN	OWN	_
PATENT API		Application Number			
FAILNIAII	LICATION	Filing Date			١
X Declaration OI		Group Art Unit			
Submitted with Initial Filing	Submitted after Initial Filing	Examiner Name			
My residence, post office address, a I believe I am the original, first and of the subject matter which is claim IMPROVED FERMEN	sole inventor (if only one name is lis ad and for which a patent is sought	sted below) or an original, first an	d joint inventor (if pl	ural names are listed below)]
OR was filed on (MM/DD/YYY Application Number I hereby state that I have reviewed a mendment specifically referred to a I acknowledge the duty to disclose in I hereby claim foreign priority benefit	and was an and was an and understand the contents of the bove	mended on (MM/DD/YYYY) above identified specification, inc	duding the claims, a	lations, § 1.56.	
I hereby claim foreign priority beneficate, or \$365(a) of any PCT in and have also identified below, by chaving a filing date before that of the			Priority	Certified Copy Attached	?
Prior Foreign Application Number(s)	Country	Foreign Filling Date (MM/DD/YYYY)	Not Claimed	YES NO	
Additional foreign application	numbers are listed on a supplemen	ital pnority sheet attached hereto	:		
I hereby claim the benefit under Tit					_
Application Number(s)	Filing Date (MM/DD/YYY)	<u>~</u> —	Additional provision application number		
60/156,791	09/30/1999	, L	are listed on a		

Burden Hour Statement: This form is estimated to take 4 hours to complete, Time will vary depending upon the needs of the inclividual case. Any comments on the amount of time you are required to complete his form should be sent to the Chef Information Officer, Patent and Trademark Office, Weshington, DC 20231, DO NOT SEND RESS OR COMPLETED PORMS TO THIS ADDRESS. SEND TO Assessart Commendations for Patents, Weshington DC 20231.

City

Indian Springs

Type a plus sign (+) inside this box •	M 6560 OS/OAPT
DECLARATION	Page 2

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §3559 of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragrept of Title 35, United States Code §121; acknowledge the duty to disclose information which is material to patentiability as defined in Title 37, Code of Federal Regulations §1 56 which became available between the filling date of the application. Parent Patent Number PCT Parent Parent Filing Date U.S. Parent (MM/DD/YYYY) (if applicable) **Application Number** Number Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto. As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith Customer Firm Name Number List Attorney(s) and/or agent(s) name and registration number below Registration Number Name Registration Name 42,516 32.891 Aaron R. Ettelman John E. Drach 36,296 18.980 Henry E. Millson, Jr. Steven J. Trzaska Additional attorney(s) and/or agent(s) named on a supplemental sheet attached hereto. OR Fill in correspondence address below Please direct all correspondence Customer 23657 x or label Name Address Address State Zip City Telephone 610-278-4929 Fax 610-278-6548 Country hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Trile 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon. A petition has been filed for this unsigned inventor Name of Sole or First Inventor: Middle W. Family Anderson Given Name Kevin Name Initial e.g. Jr. Date Inventor's Signature ОН US Citizenship US State Country Residence: City Indian Springs 4151 Weiser Court Post Office Address Post Office Address

45011

State ОН Zin

Additional inventors are being named on supplemental sheet(s) attached hereto

Country us

Page 2

Applicant

Authority

Туре а	plus :	sign (+) ins	side this box >											М 6	560	OS/C	APT
DECLARATION											ADDITIONAL INVENTOR(S) Supplemental Sheet							
Name of Additional Joint Inventor, if any:												A petition has been filed for this unsigned inventor						
Given Name		J. Douglas				Middle Initial	Fa Na		Fan	nily ne	y Wenzel				Suffix e.g. Jr.			
Invento Signat												Date						
Reside	nce: (City Cincinnati				State	OH C		C	ountry	US			Citizensh	nip	US		
Post Office Address				873 Gorham														
Post Office Address																		
City	City Cincinnati		i		State	State OH		Zip 4524		5	Cour		ry US		Applica Authori	ant ity		
Name of Additional Joint Inventor, if any: A petition has been inventor											s been fi	filed for this unsigned						
Given Name						Middle Initial		_		Fai Na	mily me					Sui e.g	fix . Jr.	
Inventor's Signature													Date					
Reside	ence:	City				State	State			С	Country				Citizens	hip		
Post C	Office .	Addre	ss															
Post C	Office	Addre	ss															
City			State				ip			Cou	Country			Applicant Authority		it y		
Name of Additional Joint Inventor, if any: A petition has been filed for this unsigned inventor																		
Given						Middle Initial					amily ame					Su e.g	uffix .g. Jr.	
Inventor's Signature														Date				
Residence: City					State	State			C	country	丄			Citizens	hip	L		
Post 6	Office	Addr	ess															
Post Office Address																		
City					Z		Zip			Country				Applic Author				
Nam	e of	Add	litio	nal Joint Ir	rvento	or, if an	y:					A p	etition h	as been t	iled for th	is u	nsigne	i
Giver Name						Middle Initial					amily ame					Suffix e.g. Jr.		
Inventor's Signature													Date					
Residence: City							State				Country				Citizen	ship		
Post	Office	Addı	ess															
Post	Office	Addı	ess															
City					State	9	T	Zip			Со	untry			Applic Autho	ant rity		
Γ	Add	dition	al inv	entors are bei	ng nam	ed on sup	pler	ner	ntal sh	eet(s) attac	hed h	ereto					